

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 21/00, C12N 1/21, 15/31, 15/63	A1	(11) International Publication Number: WO 99/46275 (43) International Publication Date: 16 September 1999 (16.09.99)
(21) International Application Number: PCT/US99/05286 (22) International Filing Date: 10 March 1999 (10.03.99) (30) Priority Data: 09/038,909 12 March 1998 (12.03.98) US (71) Applicants: SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). SMITHKLINE BEECHAM, PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors: MAY, Earl, W.; 520 Richard Road, Wayne, PA 19087 (US). EARNSHAW, David, L.; 76 St. John's Road, Stansed, Essex MC24 8JS (GB). MCDEVITT, Damien; 38 Overlook Circle, Berwyn, PA 19312 (US). (74) Agent: JACKSON, Arthur, E.; Dechert Price & Rhoads, 4000 Bell Atlantic Tower, 1717 Arch Street, Philadelphia, PA 19103-2793 (US).		(81) Designated States: JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: REPLICATIVE HELICASE dnaB OF <i>STAPHYLOCOCCUS AUREUS</i> (57) Abstract <p>The invention provides dnaB polypeptides and polynucleotides encoding dnaB polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing dnaB polypeptides to screen for antibacterial compounds.</p>		

ATTORNEY DOCKET NUMBER: 7969-088-999
SERIAL NUMBER: 09/732,091
REFERENCE: AN

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	R	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

FIELD OF THE INVENTION

10 BACKGROUND OF THE INVENTION

Clearly, there exists a need for factors, such as the dnaB embodiments of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists to find ways to prevent, ameliorate or correct such infection, dysfunction and disease.

Certain of the polypeptides of the invention possess amino acid sequence homology to a known helicases from *S. pneumoniae*, *E. coli*, *B. subtilis*, *H. pylori*, *Synechocystis* sp., *M. tuberculosis*, *S. typhimurium*, *H. influenzae* protein (see Gen Bank Accession Nos. K01174 and P37469).

5

SUMMARY OF THE INVENTION

It is an object of the invention to provide polypeptides that have been identified as *dnaB* polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2] and a known amino acid sequence or sequences of other proteins such as helicases from *S. pneumoniae*, *E. coli*, *B. subtilis*, *H. pylori*, *Synechocystis* sp., *M. tuberculosis*, *S. typhimurium*, *H. influenzae* protein.

It is a further object of the invention to provide polynucleotides that encode *dnaB* polypeptides, particularly polynucleotides that encode the polypeptide herein designated *dnaB*.

15 In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding *dnaB* polypeptides comprising a sequence set out in Table 1 [SEQ ID NO:1] which includes a full length gene, or a variant thereof.

In another particularly preferred embodiment of the invention there is a *dnaB* protein from *Staphylococcus aureus* comprising the amino acid sequence of Table 1 [SEQ ID NO:2], or a variant thereof.

20 As a further aspect of the invention there are provided isolated nucleic acid molecules encoding *dnaB*, particularly *Staphylococcus aureus* *dnaB*, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

25 In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of *dnaB* and polypeptides encoded thereby.

In another aspect of the invention there are provided polypeptides of *Staphylococcus aureus* referred to herein as *dnaB* as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

30 Among the particularly preferred embodiments of the invention are variants of *dnaB* polypeptide encoded by naturally occurring alleles of the *dnaB* gene.

In a preferred embodiment of the invention there are provided methods for producing the aforementioned dnaB polypeptides.

In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

5 In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing dnaB expression, treating disease, assaying genetic variation, and administering a dnaB polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Staphylococcus aureus* bacteria.

10 In accordance with certain preferred embodiments of this and other aspects of the invention there are provided polynucleotides that hybridize to dnaB polynucleotide sequences. particularly under stringent conditions.

In certain preferred embodiments of the invention there are provided antibodies against dnaB polypeptides.

15 In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being associated with a second
20 component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

25 In accordance with yet another aspect of the invention, there are provided dnaB agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

In a further aspect of the invention there are provided compositions comprising a dnaB polynucleotide or a dnaB polypeptide for administration to a cell or to a multicellular organism.

30 In another embodiment of the invention there is provided a computer readable medium having stored thereon a member selected from the group consisting of: a polynucleotide comprising the sequence of SEQ ID NO. 1; a polypeptide comprising the sequence of SEQ ID NO. 2; a set of polynucleotide sequences wherein at least one of said sequences comprises the

sequence of SEQ ID NO. 1; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2; a
5 polynucleotide comprising the sequence of SEQ ID NO. 1; a polypeptide comprising the sequence of SEQ ID NO. 2; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1; a data set representing a
10 polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2. A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of providing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or
15 polypeptide sequence to identify homology.

A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polypeptide sequence comprising the sequence of SEQ ID NO. 2 in a computer readable medium; and comparing said polypeptide sequence to at least one polynucleotide or polypeptide
20 sequence to identify homology.

A further embodiment of the invention provides a computer based method for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of SEQ ID NO. 1 in a computer readable medium; and screening for at least one overlapping region between said first polynucleotide sequence and a
25 second polynucleotide sequence.

A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or
30 polypeptide sequence to identify homology.

A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polypeptide sequence comprising the sequence of SEQ ID NO. 2 in a computer readable

medium; and comparing said polypeptide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

A further embodiment of the invention provides a computer based method for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of SEQ ID NO. 1 in a computer readable medium; and
5 screening for at least one overlapping region between said first polynucleotide sequence and a second polynucleotide sequence.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from
10 reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to dnaB polypeptides and polynucleotides as described in greater
15 detail below. In particular, the invention relates to polypeptides and polynucleotides of a dnaB of *Staphylococcus aureus*, which is related by amino acid sequence homology to helicases from *S. pneumoniae*, *E. coli*, *B. subtilis*, *H. pylori*, *Synechocystis* sp., *M. tuberculosis*, *S. typhimurium*, *H. influenzae* polypeptide. The invention relates especially to dnaB having the nucleotide and amino acid sequences set out in Table 1 as SEQ ID NO: 1 and SEQ ID NO: 2 respectively.

20

TABLE 1

dnaB Polynucleotide and Polypeptide Sequences

(A) Sequences from *Staphylococcus aureus* dnaB polynucleotide sequence [SEQ ID
25 NO:1].

5'-

ACGAATGTACCTGTAAATTAGATAAAGAAGTTGAAGGTACAATTCGCGTACACACAGTTGAACAATAA
AGTTGGATTGA

30

AATAAGAGGTGTAACCATTGATGGATAGAATGTATGAGCAAAATCAAATGCCACATAACAATGAAGCTG
AACAGTCTGTC

TTAGGTTCAATTATTATAGATCCAGAATTGATTAATACTACTCAGGAAGTGTTGCTTCCTGAGTCATTT
TATAGGGGTGC

CCATCAACATATTTTCCGTGCAATGATGCACTTAAATGAAGATAATAAGAAATTGATGTTGTAACATT
GATGGATCAAT

TATCGACGGAAGGTACGTTGAATGAAGCGGGTGGCCCGCAATATCTTGCAGAGTTATCTACAAATGTAC
 CAACGACGCGA
 AATGTTCAAGTATTATACTGATATCGTTTCTAAGCATGCATTAAAACGTAGATTGATTCAAACGAGAT
 AGTATTGCCAA
 5 TGATGGATATAATGATGAACCTGAACCTAGATGCGATTTTAAGTGATGCAGAACGTCGAATTTTAGAGCT
 ATCATCTTCTC
 GTGAAAGCGATGGCTTTAAAGACATTCGAGATGTCTTAGGACAAGTATATGAAACAGCTGAAGAGCTTG
 ATCAAAATAGT
 GGTCAAACACCAGGTATTCCTACAGGATATCGAGATTTAGACCAAATGACAGCAGGGTTCAACCGAAAT
 10 GATTTAATTAT
 CCTTGCAGCGCGTCCATCTGTAGGTAAGACTGCGTTGCGACTTAATATTGCACAAAAAGTTGCAACGCA
 TGAAGATATGT
 ATACAGTTGGTATTTTCTCACTAGAGATGGGTGCTGATCAGTTAGCCACACGTATGATTTGTAGTTCTG
 GTAATGTTGAC
 15 TCAAACCGCTTAAGAACCGGTACTATGACTGAGGAAGATTGGAGTCGTTTTACTATAGCGGTAGGTAAA
 TTATCACGTAC
 GAAGATTTTTATTGATGATACACCGGGCATTGGAATTAATGATCTACGTTCTAAATGTCGTCGATTAAA
 GCAAGAACATG
 GCTTAGACATGATTGTAATTGACTACTTACAGTTGATTCAAGGTAGTGGTTCACGTGCGTCCGATAACA
 20 GACAACAGGAA
 GTTTCTGAAATCTCTCGTACATTAAAAGCATTAGCTCGTGAATTAGAATGTCCAGTTATCGCGTTGAGT
 CAGTTATCTCG
 TGGTGTGAACAACGACAAGATAAACGTCCAATGATGAGTGATATTCGTGAATCTGGTTCGATTGAGCA
 AGATGCCGATA
 25 TCGTTGCATTCTTATACCGTGATGATTACTATAACCGTGGCGGCGATGAAGATGATGACGATGATAGTG
 GTTTTGAGCCA
 CAAACGAATGATGAAAACGGTGAAATTGAAATCATCATTGCTAAGCAACGTAATGGTCCAACAGGCACA
 GTTAAGTTACA
 CTTTATGAAACAATATAATAAATTTACAGATATCGATTATGCACATGCTGATATGATGTAAAAAGTTT
 30 TTCCGTACAAT
 AATCATCTCGATGATAAAATTGTACGGTTTTTATTCGTTCTGAACGGGTTGATATATGTTAAGTTTGT
 GTATTGAAGT
 G-3'

35 (B) *Staphylococcus aureus* dnaB polypeptide sequence deduced from the polynucleotide
 sequence in this table [SEQ ID NO:2].

NH₂-

MDRMYEQNQMPHNNEAEQSVLGSIIIDPELINTTQEVLLPESFYRGHQHIFRAMMHLNEDNKEIDVVT
 LMDQLSTEGTL

NEAGGPQYLAELSTNVPTTRNVQYYTDIVSKHALKRRLIQTADSIANDGYNDELELDAILSDAERRILE
 LSSSRESDFK
 DIRDLVGLQVYETAELDQNSGQTPGIPTGYRDLDMQTAGFNRNDLIILAARPSVGKTAFALNIAQKVAT
 HEDMYTVGIFS
 5 LEMGADQLATRMICSSGNVDSNRLRTGTMTEEDWSRFTIAVGKLSRTKIFIDDTGIRINDLRSKCRL
 KQEHGLDMIVI
 DYQLIQGSGSRASDNRRQEVSEISRTLKALARELECPVIALSQLSRGVEQRQDKRPMMSDIRESGSIE
 QDADIVAFLYR
 DDYNRGGDEDDDDSGFEPQTNDENGEIEIIIAKQRNGPTGTVKLHFMKQYNKFTDIDYAHADMM. -
 10 COOH

(C) Polynucleotide sequences comprising *Staphylococcus aureus* dnaB ORF sequence
 [SEQ ID NO:3].

5' -ATCCCTAGGATATACGAATGTNCCTGTAAATTAGATAAAGAAGTTGAAGGTACAATTCGCGTACA
 15 CACAGTTGANCAATAAAGTTGGATTGAAATAAGAGGTGTAACCATTGATGATAGAATGTATGAGC
 AAAATCAANTGCCACATAACAATGAAGCTGAACAGTCTGTCTTAGGTTCAANTATTATAGATCCAGA
 ATTGATTAATACTACTCAGGAAGTTTNGCTTCCTGAGTCGTTTTATAGGGGTGCCCATCAACATATT
 TTCCGTGCAATGATGCACTTAAATGAAGATAATAAAGAAATTGATGTTGTAACATTGATGGATCAAT
 TATCGACGGAAGGTACGTTGAATGAAGCGGGTGGCCCGCAATATCTTGAGAGTTATCTACAAATGT
 20 NCCAACGACGCGAAATGTCCAGTATTATACTGATATCGNCTCTTANGCATGCATTAAANACGTAGGA
 TTGATTCAAATGCAGATAGTATTGCCAATGATGGATATANTGATGAACNNGGACTAGATGCGATTT
 AAGNGATGCAGAACGTCGNATTTTAGAG-3'

(D) *Staphylococcus aureus* dnaB polypeptide sequence deduced from the polynucleotide
 25 ORF sequence in this table [SEQ ID NO:4].

NH₂-
 PDIRMXLLNIKKLVQFAYTQLXNKVGLKEVPFMDRMYEQNQXPHNNEAEQSVLGSXIIDPELINTTQE
 VXLPESEFYRGAHQHIFRAMMHLNEDNKEIDVVTLMQDQSTEGTLNEAGGPQYLAELSTNVPTTRNVQYY
 TDIXSXACIKXVGLIQTADSIANDGYXDELGLDAIXMQNVXF-COOH
 30

Deposited materials

A deposit containing a *Staphylococcus aureus* WCUH 29 strain has been deposited with
 the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar
 Drive, Aberdeen AB2 1RY, Scotland on 11 September 1995 and assigned NCIMB Deposit No.
 35 40771, and referred to as *Staphylococcus aureus* WCUH29 on deposit. The *Staphylococcus aureus*
 strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains the full length dnaB gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

5 The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required
10 under 35 U.S.C. §112.


A license may be required to make, use or sell the deposited strain, and compounds derived therefrom, and no such license is hereby granted.

One aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH 29 strain contained in the
15 deposited strain. Further provided by the invention are dnaB nucleotide sequences of the DNA in the deposited strain and amino acid sequences encoded thereby. Also provided by the invention are dnaB polypeptide sequences isolated from the deposited strain and amino acid sequences derived therefrom.

Polypeptides

20 The polypeptides of the invention include a polypeptide of Table 1 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of dnaB, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NO:1] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NO:2] and more preferably at least 90% identity to a polypeptide of Table 1
25 [SEQ ID NO:2] and still more preferably at least 95% identity to a polypeptide of Table 1 [SEQ ID NO:2] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

The invention also includes polypeptides of the formula:



30 wherein, at the amino terminus, X is hydrogen or a metal, and at the carboxyl terminus, Y is hydrogen or a metal, R₁ and R₃ are any amino acid residue, m is an integer between 1 and 1000 or zero, n is an integer between 1 and 1000 or zero, and R₂ is an amino acid sequence of the invention, particularly an amino acid sequence selected from Table 1. In the formula above R₂ is oriented so

that its amino terminal residue is at the left, bound to R_1 , and its carboxy terminal residue is at the right, bound to R_3 . Any stretch of amino acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

5 A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with dnaB polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

10 Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NO:2], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Staphylococcus aureus*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and
15 coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Also preferred are biologically active fragments which are those fragments that mediate activities of dnaB, including those with a similar activity or an improved activity, or with a decreased
20 undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Staphylococcus aureus* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

Variants that are fragments of the polypeptides of the invention may be employed for
25 producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

In addition to the standard single and triple letter representations for amino acids, the term "X" or "Xaa" may also be used in describing certain polypeptides of the invention. "X" and "Xaa" mean that any of the twenty naturally occurring amino acids may appear at such a
30 designated position in the polypeptide sequence.

Polynucleotides

Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the dnaB polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2] and polynucleotides closely related thereto and variants thereof.

Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NO:1], a polynucleotide of the invention encoding dnaB polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Staphylococcus aureus* WCUH 29 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NO:1], typically a library of clones of chromosomal DNA of *Staphylococcus aureus* WCUH 29 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING. A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1] was discovered in a DNA library derived from *Staphylococcus aureus* WCUH 29.

The DNA sequence set out in Table 1 [SEQ ID NO:1] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 101 and the stop codon which begins at nucleotide number 1499 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.


The dnaB protein of the invention is structurally related to other proteins of the Replicative Helicase family.

The invention provides a polynucleotide sequence identical over its entire length to a coding sequence in Table 1 [SEQ ID NO:1]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature

polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 101 to the nucleotide immediately upstream of or including nucleotide 1499 set forth in SEQ ID NO:1 of Table 1, both of which encode the dnaB polypeptide.

The invention also includes polynucleotides of the formula:



wherein, at the 5' end of the molecule, X is hydrogen or a metal or together with Y defines a covalent bond, and at the 3' end of the molecule, Y is hydrogen or a metal or together with X defines the covalent bond, each occurrence of R_1 and R_3 is independently any nucleic acid residue. m is an integer between 1 and 3000 or zero, n is an integer between 1 and 3000 or zero, and R_2 is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from Table 1. In the polynucleotide formula above R_2 is oriented so that its 5' end residue is at the left, bound to R_1 , and its 3' end residue is at the right, bound to R_3 . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. Where, in a preferred embodiment, X and Y together define a covalent bond, the polynucleotide of the above formula is a closed, circular polynucleotide, which can be a double-stranded polynucleotide wherein the formula shows a first strand to which the second strand is complementary. In another preferred embodiment m and/or n is an integer between 1 and 1000.

It is most preferred that the polynucleotides of the inventions are derived from *Staphylococcus aureus*, however, they may preferably be obtained from organisms of the same

taxonomic genus. They may also be obtained, for example, from organisms of the same taxonomic family or order.

5 The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Staphylococcus aureus* dnaB having an amino acid sequence set out in Table 1 [SEQ ID NO:2]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

10 The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

15 Further particularly preferred embodiments are polynucleotides encoding dnaB variants, that have the amino acid sequence of dnaB polypeptide of Table 1 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of dnaB.

20 Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding dnaB polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NO:2], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding dnaB polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over
25 their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

30 Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of Table 1 [SEQ ID NO:1].

The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under

stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising:

5 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y.,

10 (1989), particularly Chapter 11 therein.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or

15 a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding dnaB

20 and to isolate cDNA and genomic clones of other genes that have a high sequence identity to the dnaB gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

For example, the coding region of the dnaB gene may be isolated by screening using a DNA

25 sequence provided in Table 1 [SEQ ID NO: 1] to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The polynucleotides and polypeptides of the invention may be employed, for example, as

30 research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

Polynucleotides of the invention that are oligonucleotides derived from the sequences of Table 1 [SEQ ID NOS:1 or 2] may be used in the processes herein as described, but preferably

for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

5 The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino
10 acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

15 In addition to the standard A, G, C, T/U representations for nucleic acid bases, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA bases may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a base that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the
20 effect of generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more
25 prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

The invention further comprises an isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:4. The invention further comprises an isolated polynucleotide comprising a polynucleotide having at least a 70% identity to the polynucleotide sequence of SEQ ID NO:3.

Vectors, host cells, expression

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production

of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING. A LABORATORY MANUAL*, (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic, Prognostic, Serotyping and Mutation Assays

This invention is also related to the use of the *dnaB* polynucleotides of the invention for use as diagnostic reagents. Detection of *dnaB* in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the *dnaB* gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior to analysis. RNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled *dnaB* polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

Cells carrying mutations or polymorphisms (allelic variations) in the gene of the invention may also be detected at the DNA or RNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to used RT-PCR in conjunction with automated detection systems, such as, for example,


GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding dnaB can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

Table 2

Primers for amplification of dnaB polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
5	5'-AAATTAGATAAAGAAGTTGAAGGT-3'
10 6	5'-CGAGATGATTATTGTACGAAAAA-3'

The invention also includes primers of the formula:



wherein, at the 5' end of the molecule, X is hydrogen or a metal, and at the 3' end of the molecule, Y is hydrogen or a metal, R₁ and R₃ is any nucleic acid residue, m is an integer between 1 and 20 or zero, n is an integer between 1 and 20 or zero, and R₂ is a primer sequence of the invention, particularly a primer sequence selected from Table 2. In the polynucleotide formula above R₂ is oriented so that its 5' end residue is at the left, bound to R₁, and its 3' end residue is at the right, bound to R₃. Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer being complementary to a region of a polynucleotide of Table 1. In a preferred embodiment m and/or n is an integer between 1 and 10.

The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying dnaB DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Staphylococcus aureus*, comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having a sequence of Table 1 [SEQ ID NO: 1]. Increased or decreased expression of dnaB polynucleotide can be measured using any one of the methods well known in the art for the

quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of dnaB protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a dnaB protein, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

The polynucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual microbial chromosome, particularly a *Staphylococcus aureus* chromosome. The mapping of relevant sequences to a chromosome according to the present invention is an important first step in correlating those sequences with gene associated with microbial pathogenicity and disease, or to chromosomal regions critical to the growth, survival and/or ecological niche. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, microbial genomic sequences available on the World Wide Web. The relationship between genes and microbial pathogenicity, disease, or to genome regions critical to the growth, survival and/or ecological niche that have been mapped to the same chromosomal region are then identified using methods to define a genetic relationship between the gene and another gene or phenotype, such as by linkage analysis (coinheritance of physically adjacent genes).

The differences in the RNA or genomic sequence between microbes of differing phenotypes can also be determined. If a mutation or sequence is observed in some or all of the microbes of a certain phenotype, but not in any microbes lacking that phenotype, then the mutation or sequence is likely to be the causative agent of the phenotype. In this way, chromosomal regions may be identified that confer microbial pathogenicity, growth characteristics, survival characteristics and/or ecological niche characteristics.

Differential Expression

The polynucleotides and polynucleotides of the invention may be used as reagents for differential screening methods. There are many differential screening and differential display methods known in the art in which the polynucleotides and polypeptides of the invention may be used. For example, the differential display technique is described by Chuang *et al.*, *J. Bacteriol.* 175:2026-2036 (1993). This method identifies those genes which are expressed in

an organism by identifying mRNA present using randomly-primed RT-PCR. By comparing pre-infection and post infection profiles, genes up and down regulated during infection can be identified and the RT-PCR product sequenced and matched to ORF 'unknowns'.

In Vivo Expression Technology (IVET) is described by Camilli *et al.*, *Proc. Nat'l. Acad. Sci. USA*, 91:2634-2638 (1994). IVET identifies genes up-regulated during infection when compared to laboratory cultivation, implying an important role in infection. ORF identified by this technique are implied to have a significant role in infection establishment and/or maintenance. In this technique random chromosomal fragments of target organism are cloned upstream of a promoter-less recombinase gene in a plasmid vector. This construct is introduced into the target organism which carries an antibiotic resistance gene flanked by resolvase sites. Growth in the presence of the antibiotic removes from the population those fragments cloned into the plasmid vector capable of supporting transcription of the recombinase gene and therefore have caused loss of antibiotic resistance. The resistant pool is introduced into a host and at various times after infection bacteria may be recovered and assessed for the presence of antibiotic resistance. The chromosomal fragment carried by each antibiotic sensitive bacterium should carry a promoter or portion of a gene normally upregulated during infection. Sequencing upstream of the recombinase gene allows identification of the up regulated gene.

RT-PCR may also be used to analyze gene expression patterns. For RT PCR using the polynucleotides of the invention, messenger RNA is isolated from bacterial infected tissue, e.g., 48 hour murine lung infections, and the amount of each mRNA species assessed by reverse transcription of the RNA sample primed with random hexanucleotides followed by PCR with gene specific primer pairs. The determination of the presence and amount of a particular mRNA species by quantification of the resultant PCR product provides information on the bacterial genes which are transcribed in the infected tissue. Analysis of gene transcription can be carried out at different times of infection to gain a detailed knowledge of gene regulation in bacterial pathogenesis allowing for a clearer understanding of which gene products represent targets for screens for antibacterials. Because of the gene specific nature of the PCR primers employed it should be understood that the bacterial mRNA preparation need not be free of mammalian RNA. This allows the investigator to carry out a simple and quick RNA preparation from infected tissue to obtain bacterial mRNA species which are very short lived in the bacterium (in the order of 2 minute half-lives). Optimally the bacterial mRNA is prepared from infected murine lung tissue by mechanical disruption in the presence of TRIzol

(GIBCO-BRL) for very short periods of time, subsequent processing according to the manufacturers of TRIzol reagent and DNAase treatment to remove contaminating DNA. Preferably the process is optimised by finding those conditions which give a maximum amount of *Staphylococcus aureus* 16S ribosomal RNA as detected by probing Northern blots with a suitably labelled sequence specific oligonucleotide probe. Typically a 5' dye labelled primer is used in each PCR primer pair in a PCR reaction which is terminated optimally between 8 and 25 cycles. The PCR products are separated on 6% polyacrylamide gels with detection and quantification using GeneScanner (manufactured by ABI).

Each of these techniques may have advantages or disadvantage depending on the particular application. The skilled artisan would choose the approach that is the most relevant with the particular end use in mind.

Antibodies

The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies.

Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-dnaB or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides to purify the polypeptides by affinity chromatography.

5 Thus, among others, antibodies against dnaB-polypeptide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide according to the invention, interfere with the immediate physical interaction between pathogen and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

15 The polypeptide, such as an antigenically or immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

25 Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized"; where the complementarity determining region(s) of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

30 The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff *et al.*, *Hum Mol Genet* 1992, 1:363, Manthorpe *et al.*, *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu *et al.*, *J Biol Chem.* 1989: 264.16985).

coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef. *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al. *Science* 1989:243,375), particle bombardment (Tang et al. *Nature* 1992, 356:152, Eisenbraun et al. *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

Antagonists and agonists - assays and molecules

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of dnaB polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising dnaB polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a dnaB agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the dnaB polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, i.e., without inducing the effects of dnaB polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in dnaB polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for dnaB antagonists is a competitive assay that combines dnaB and a potential antagonist with dnaB-binding molecules, recombinant dnaB binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. The dnaB molecule can be labeled, such as by radioactivity or a colorimetric compound, such that the number of dnaB molecules bound to a binding molecule or

converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing dnaB-induced activities, thereby preventing the action of dnaB by excluding dnaB from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of dnaB.

Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block dnaB protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial dnaB proteins that mediate tissue damage and; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat diseases.

Helicobacter pylori (herein *H. pylori*) bacteria infect the stomachs of over one-third of the world's population causing stomach cancer, ulcers, and gastritis (International Agency for Research on Cancer (1994) Schistosomes, Liver Flukes and *Helicobacter Pylori* (International Agency for Research on Cancer, Lyon, France; <http://www.uicc.ch/ecp/ecp2904.htm>). Moreover, the international Agency for Research on Cancer recently recognized a cause-and-effect relationship between *H. pylori* and gastric adenocarcinoma, classifying the bacterium as a Group I (definite) carcinogen. Preferred antimicrobial compounds of the invention (agonists and antagonists of *dnaB*) found using screens provided by the invention, particularly broad-spectrum antibiotics, should be useful in the treatment of *H. pylori* infection. Such treatment should decrease the advent of *H. pylori*-induced cancers, such as gastrointestinal carcinoma. Such treatment should also cure gastric ulcers and gastritis.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with *dnaB*, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Staphylococcus aureus* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of *dnaB*, or a fragment or a variant thereof, for expressing *dnaB*, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a *dnaB* or protein coded therefrom, wherein the composition comprises a recombinant *dnaB* or protein coded therefrom comprising

DNA which codes for and expresses an antigen of said dnaB or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

5 A dnaB polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large
10 co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and
15 immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in
20 animal models of infection with *Staphylococcus aureus* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic treatments of
25 bacterial infection, particularly *Staphylococcus aureus* infection, in mammals, particularly humans.

The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or
30 connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations
5 suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example,
10 sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

15 While the invention has been described with reference to certain dnaB protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

Compositions, kits and administration

20 The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the
25 invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

30 Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual
5 as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application
for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops,
mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate
conventional additives, including, for example, preservatives, solvents to assist drug
10 penetration, and emollients in ointments and creams. Such topical formulations may also
contain compatible conventional carriers, for example cream or ointment bases, and ethanol or
oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight
of the formulation; more usually they will constitute up to about 80% by weight of the
formulation.

15 For administration to mammals, and particularly humans, it is expected that the daily
dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1
mg/kg. The physician in any event will determine the actual dosage which will be most suitable
for an individual and will vary with the age, weight and response of the particular individual.
The above dosages are exemplary of the average case. There can, of course, be individual
20 instances where higher or lower dosage ranges are merited, and such are within the scope of
this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e.,
devices that are introduced to the body of an individual and remain in position for an extended
time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular
25 grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory
peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a
systemic effect against relevant bacteria shortly before insertion of an in-dwelling device.
Treatment may be continued after surgery during the in-body time of the device. In addition,
30 the composition could also be used to broaden perioperative cover for any surgical technique to
prevent bacterial wound infections, especially *Staphylococcus aureus* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be
considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia.

Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis. Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Sequence Databases and Algorithms

The polynucleotide and polypeptide sequences of the invention are particularly useful as components in databases useful for search analyses as well as in sequence analysis algorithms. As used in this section entitled Databases and Algorithms and in claims related thereto, the terms "polynucleotide of the invention" and "polynucleotide sequence of the invention" mean any detectable chemical or physical characteristic of a polynucleotide of the invention that is or may be reduced to or stored in a computer readable form. For example, chromatographic scan data or peak data, photographic data or scan data therefrom, called bases, and mass spectrographic data. As used in this section entitled Databases and Algorithms and in claims related thereto, the terms "polypeptide of the invention" and "polypeptide sequence of the invention" mean any detectable chemical or physical characteristic of a polypeptide of the invention that is or may be reduced to or stored in a computer readable form. For example, chromatographic scan data or peak data, photographic data or scan data therefrom, and mass spectrographic data.

The invention provides computer readable medium having stored thereon sequences of the invention. For example, a computer readable medium is provided having stored thereon a member selected from the group consisting of: a polynucleotide comprising the sequence of a

polynucleotide of the invention; a polypeptide comprising the sequence of a polypeptide sequence of the invention; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of a polynucleotide sequence of the invention; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of a polypeptide sequence of the invention; a data set representing a polynucleotide sequence comprising the sequence of polynucleotide sequence of the invention; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of a polypeptide sequence of the invention; a polynucleotide comprising the sequence of a polynucleotide sequence of the invention; a polypeptide comprising the sequence of a polypeptide sequence of the invention; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of a polynucleotide sequence of the invention; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of a polypeptide sequence of the invention; a data set representing a polynucleotide sequence comprising the sequence of a polynucleotide sequence of the invention; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of a polypeptide sequence of the invention. The computer readable medium can be any composition of matter used to store information or data, including, for example, commercially available floppy disks, tapes, hard drives, compact disks, and video disks.

Also provided by the invention are methods for the analysis of character sequences, particularly genetic sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, RNA structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, and sequencing chromatogram peak analysis.

A computer based method is provided for performing homology identification. This method comprises the steps of providing a polynucleotide sequence comprising the sequence a polynucleotide of the invention in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said polypeptide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

A computer based method is still further provided for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and screening for at least one overlapping region between said first polynucleotide sequence and a second polynucleotide sequence.

A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said polypeptide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

A further embodiment of the invention provides a computer based method for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and screening for at least one overlapping region between said first polynucleotide sequence and a second polynucleotide sequence.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety.

GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Disease(s)" means and disease caused by or related to infection by a bacteria, including disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous

abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

- 5 "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M.,
10 ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and
15 *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.* 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include,
20 but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman
25 algorithm may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

- 1) Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)
Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci. USA.* 89:10915-10919 (1992)
30 Gap Penalty: 12
Gap Length Penalty: 4

A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

- 5 1) Algorithm: Needleman and Wunsch, J. Mol Biol. 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

- Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the
10 default parameters for nucleic acid comparisons.

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

- (1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100%
15 identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO: 1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations
20 may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and
25 then subtracting that product from said total number of nucleotides in SEQ ID NO:1. or:

$$n_n \leq x_n - (x_n \cdot y),$$

- wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ
30 ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence

encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

By way of example, a polynucleotide sequence of the present invention may be
 5 identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal
 10 positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then
 15 subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_n \leq x_n - (x_n \cdot y),$$

wherein n_n is the number of amino acid alterations, x_n is the total number of amino acids in
 20 SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50,60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide
 25 reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO: 2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may
 30 occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the

total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$5 \quad n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2. or:

25

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

"Isolated" means altered "by the hand of man" from its natural state, i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a

polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxiribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and

other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Scifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally,

differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1 Strain selection, Library Production and Sequencing

The polynucleotide having a DNA sequence given in Table 1 [SEQ ID NO:1] was obtained from a library of clones of chromosomal DNA of *Staphylococcus aureus* in *E. coli*. The sequencing data from two or more clones containing overlapping *Staphylococcus aureus* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1 and 2 below.

Total cellular DNA is isolated from *Staphylococcus aureus* WCUH 29 according to standard procedures and size-fractionated by either of two methods.

Method 1

Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI. the library packaged by standard procedures and *E. coli* infected with the packaged library. The library is amplified by standard procedures.

Method 2

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., *RsaI*, *Pall*, *AluI*, *BshI235I*), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the

vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Example 2 dnaB Characterization

5 S. aureus DnaB expression:

Using PCR amplification from *S. aureus* WCUH29 genomic DNA (5' primer: AGCATCATCCCATGGATAGAATGTATGAGCAA, [SEQ ID NO: 7] 3' primer: ATCTATTCCTCGAGCATCATATCAGCATGTGCAT [SEQ ID NO: 8]), the DnaB gene was isolated and subcloned (NcoI/XhoI) into a suitable protein expression vector, pBluepET28a, made in house from pBluescript (Stratagene) and pET28a (Invitrogen), linked to a C-terminal polyhistidine affinity tag. Protein expression was carried out in the *E. coli* strain LW29(DE3) containing pRI952 which encodes rare *E. coli* tRNAs for Ile and Arg. Induction with 0.3 mM IPTG at 37C for 3hrs yielded a 53kD MW product on a 4-12% NuPAGE gel (Novex), consistent with the expected size of the protein fusion. The protein was insoluble when induced at 37C, solubility improved when induction temperature was lowered to 24C.

10

15

What is claimed is:

1. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
2. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the *dnaB* gene contained in the *Staphylococcus aureus*.
3. An isolated polynucleotide comprising a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
4. An isolated polynucleotide that is complementary to the polynucleotide of claim 1
5. The polynucleotide of Claim 1 wherein the polynucleotide is DNA or RNA
6. The polynucleotide of Claim 1 comprising the nucleic acid sequence set forth in SEQ ID NO:1.
7. The polynucleotide of Claim 1 comprising nucleotide 101 to the stop codon which begins at nucleotide number 1499 set forth in SEQ ID NO:1.
8. The polynucleotide of Claim 1 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
9. A vector comprising the polynucleotide of Claim 1.
10. A host cell comprising the vector of Claim 9.
11. A process for producing a polypeptide comprising: expressing from the host cell of Claim 10 a polypeptide encoded by said DNA.
12. A process for producing a *dnaB* polypeptide or fragment comprising culturing a host of claim 10 under conditions sufficient for the production of said polypeptide or fragment.
13. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
14. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.
15. An antibody against the polypeptide of claim 14.
16. An antagonist which inhibits the activity or expression of the polypeptide of claim 14.

17. A method for the treatment of an individual in need of dnaB polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 14.

18. A method for the treatment of an individual having need to inhibit dnaB polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.

19. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 14 in an individual comprising:

- (a) determining a nucleic acid sequence encoding said polypeptide, and/or
- (b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.

20. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 14 comprising:

contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;

and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.

21. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with dnaB polypeptide of claim 14, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.

22. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of dnaB polypeptide of claim 14, or fragment or a variant thereof, for expressing said dnaB polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.

23. A computer readable medium having stored thereon a member selected from the group consisting of: a polynucleotide comprising the sequence of SEQ ID NO. 1; a polypeptide comprising the sequence of SEQ ID NO. 2; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1; a set of

polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2; a polynucleotide comprising the sequence of SEQ ID NO. 1; a polypeptide comprising the sequence of SEQ ID NO. 2; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2.

24. A computer based method for performing homology identification, said method comprising the steps of providing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

25. A further embodiment of the invention provides a computer based method for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of SEQ ID NO. 1 in a computer readable medium; and screening for at least one overlapping region between said first polynucleotide sequence and a second polynucleotide sequence.

26. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:4.

27. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to the polynucleotide sequence of SEQ ID NO:3.

SEQUENCE LISTING

<110> SmithKline Beecham Corporation
SmithKline Beecham plc

<120> dnaB

<130> GM10133

<150> 09/038,909

<151> 1998-03-12

<160> 8

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1601

<212> DNA

<213> Staphylococcus aureus

<400> 1

acgaatgtac ctgttaaatt agataaagaa gttgaaggta caattcgcgt acacacagtt
60
gaacaataaa gttggattga aataagaggt gtaaccattc atggatagaa tgtatgagca
120
aatcaaatg ccacataaca atgaagctga acagtctgtc ttaggttcaa ttattataga
180
tccagaattg attaatacta ctgaggaagt gttgcttct gagtcatttt ataggggtgc
240
ccatcaacat attttccgtg caatgatgca cttaaataag gataataaag aaattgatgt
300
tgtaacattg atggatcaat tatcgacgga aggtacgttg aatgaagcgg gtggcccgca
360
atatcttgca gagttatcta caaatgtacc aacgacgca aatgttcagt attatactga
420
tatcgtttct aagcatgcat taaaacgtag attgattcaa actgcagata gtattgccaa
480
tgatggatat aatgatgaac ttgaactaga tgcgatttta agtgatgcag aacgtcgaat
540

tttagagcta tcattctctc gtgaaagcga tggctttaaa gacattcgag atgtcttagg
600
acaagtatat gaaacagctg aagagcttga tcaaaatagt ggtcaaacac caggatttcc
660
tacaggatat cgagatttag accaaatgac agcaggggtc aaccgaaatg atttaattat
720
ccttgacgag cgtccatctg taggtaagac tgcgttcgca cttaatattg cacaaaaagt
780
tgcaacgcat gaagatatgt atacagttgg tattttctca ctagagatgg gtgctgatca
840
gtagccaca cgtatgattt gtagttctgg taatgttgac tcaaaccgct taagaaccgg
900
tactatgact gaggaagatt ggagtcgttt tactatagcg gtaggtaaata tatcacgtac
960
gaagattttt attgatgata caccgggcat tcgaattaat gatctacgtt ctaaagtctg
1020
tcgattaaag caagaacatg gcttagacat gattgtaatt gactacttac agttgattca
1080
aggtagtggc tcacgtgcgt ccgataacag acaacaggaa gtttctgaaa tctctcgtac
1140
attaaaagca ttagctcgtg aattagaatg tccagttatc gcgttgagtc agttatctcg
1200
tggtgttgaa caacgacaag ataaacgtcc aatgatgagt gatattcgtg aatctgggtc
1260
gattgagcaa gatgccgata tcgttgcatc cttataccgt gatgattact ataaccgtgg
1320
cggcgatgaa gatgatgacg atgatagtgg ttttgagcca caaacgaatg atgaaaacgg
1380
tgaaattgaa atcatcattg ctaagcaacg taatggtcca acaggcacag ttaagttaca
1440
ctttatgaaa caatataata aatttacaga tatcgattat gcacatgctg atatgatgta
1500
aaaaagtttt tccgtacaat aatcatctcg atgataaaat tgtacggttt ttatttcggt
1560
ctgaacgggt tgatatatgt taagtttgtg tatttgaagt g
1601

<210> 2

<211> 466

<212> PRT

<213> Staphylococcus aureus

<400> 2

```

Met Asp Arg Met Tyr Glu Gln Asn Gln Met Pro His Asn Asn Glu Ala
 1           5           10           15
Glu Gln Ser Val Leu Gly Ser Ile Ile Ile Asp Pro Glu Leu Ile Asn
      20           25           30
Thr Thr Gln Glu Val Leu Leu Pro Glu Ser Phe Tyr Arg Gly Ala His
      35           40           45
Gln His Ile Phe Arg Ala Met Met His Leu Asn Glu Asp Asn Lys Glu
      50           55           60
Ile Asp Val Val Thr Leu Met Asp Gln Leu Ser Thr Glu Gly Thr Leu
      65           70           75           80
Asn Glu Ala Gly Gly Pro Gln Tyr Leu Ala Glu Leu Ser Thr Asn Val
      85           90           95
Pro Thr Thr Arg Asn Val Gln Tyr Tyr Thr Asp Ile Val Ser Lys His
      100           105           110
Ala Leu Lys Arg Arg Leu Ile Gln Thr Ala Asp Ser Ile Ala Asn Asp
      115           120           125
Gly Tyr Asn Asp Glu Leu Glu Leu Asp Ala Ile Leu Ser Asp Ala Glu
      130           135           140
Arg Arg Ile Leu Glu Leu Ser Ser Ser Arg Glu Ser Asp Gly Phe Lys
      145           150           155           160
Asp Ile Arg Asp Val Leu Gly Gln Val Tyr Glu Thr Ala Glu Glu Leu
      165           170           175
Asp Gln Asn Ser Gly Gln Thr Pro Gly Ile Pro Thr Gly Tyr Arg Asp
      180           185           190
Leu Asp Gln Met Thr Ala Gly Phe Asn Arg Asn Asp Leu Ile Ile Leu
      195           200           205
Ala Ala Arg Pro Ser Val Gly Lys Thr Ala Phe Ala Leu Asn Ile Ala
      210           215           220
Gln Lys Val Ala Thr His Glu Asp Met Tyr Thr Val Gly Ile Phe Ser
      225           230           235           240
Leu Glu Met Gly Ala Asp Gln Leu Ala Thr Arg Met Ile Cys Ser Ser
      245           250           255
Gly Asn Val Asp Ser Asn Arg Leu Arg Thr Gly Thr Met Thr Glu Glu
      260           265           270
Asp Trp Ser Arg Phe Thr Ile Ala Val Gly Lys Leu Ser Arg Thr Lys
      275           280           285
Ile Phe Ile Asp Asp Thr Pro Gly Ile Arg Ile Asn Asp Leu Arg Ser
      290           295           300
Lys Cys Arg Arg Leu Lys Gln Glu His Gly Leu Asp Met Ile Val Ile
      305           310           315           320
Asp Tyr Leu Gln Leu Ile Gln Gly Ser Gly Ser Arg Ala Ser Asp Asn

```

325 330 335
 Arg Gln Gln Glu Val Ser Glu Ile Ser Arg Thr Leu Lys Ala Leu Ala
 340 345 350
 Arg Glu Leu Glu Cys Pro Val Ile Ala Leu Ser Gln Leu Ser Arg Gly
 355 360 365
 Val Glu Gln Arg Gln Asp Lys Arg Pro Met Met Ser Asp Ile Arg Glu
 370 375 380
 Ser Gly Ser Ile Glu Gln Asp Ala Asp Ile Val Ala Phe Leu Tyr Arg
 385 390 395 400
 Asp Asp Tyr Tyr Asn Arg Gly Gly Asp Glu Asp Asp Asp Asp Ser
 405 410 415
 Gly Phe Glu Pro Gln Thr Asn Asp Glu Asn Gly Glu Ile Glu Ile Ile
 420 425 430
 Ile Ala Lys Gln Arg Asn Gly Pro Thr Gly Thr Val Lys Leu His Phe
 435 440 445
 Met Lys Gln Tyr Asn Lys Phe Thr Asp Ile Asp Tyr Ala His Ala Asp
 450 455 460
 Met Met
 465

<210> 3
 <211> 562
 <212> DNA
 <213> Staphylococcus aureus

<220>
 <221> misc_feature
 <222> (1)...(562)
 <223> n = A,T,C or G

<400> 3
 atccctagga tatacgaatg tncctgttaa attagataaa gaagttgaag gtacaattcg
 60
 cgtacacaca gttgancaat aaagttggat tgaaataaga ggtgtaacca ttcattgata
 120
 gaatgtatga gcaaaatcaa ntgccacata acaatgaagc tgaacagtct gtcttaggtt
 180
 canttattat agatccagaa ttgattaata ctactcagga agtttngctt cctgagtcgt
 240
 tttatagggg tgcccatcaa catattttcc gtgcaatgat gcacttaaata gaagataata
 300

aagaaattga tgttgtaaca ttgatggatc aattatcgac ggaaggtacg ttgaatgaag
 360
 cgggtggccc gcaatatctt gcagagttat ctacaaatgt nccaacgacg cgaaatgtcc
 420
 agtattatac tgatatcgnc tcttangcat gcattaaana cgtaggattg attcaaactg
 480
 cagatagtat tgccaatgat ggatatantg atgaactngg actagatgcg atttaagnga
 540
 tgcagaacgt cgnatttttag ag
 562

<210> 4
 <211> 180
 <212> PRT
 <213> Staphylococcus aureus

 <220>
 <221> VARIANT
 <222> (1)...(180)
 <223> Xaa = Any Amino Acid

<400> 4
 Pro Asp Ile Arg Met Xaa Leu Leu Asn Ile Lys Lys Leu Lys Val Gln
 1 5 10 15
 Phe Ala Tyr Thr Gln Leu Xaa Asn Lys Val Gly Leu Lys Glu Val Pro
 20 25 30
 Phe Met Asp Arg Met Tyr Glu Gln Asn Gln Xaa Pro His Asn Asn Glu
 35 40 45
 Ala Glu Gln Ser Val Leu Gly Ser Xaa Ile Ile Asp Pro Glu Leu Ile
 50 55 60
 Asn Thr Thr Gln Glu Val Xaa Leu Pro Glu Ser Phe Tyr Arg Gly Ala
 65 70 75 80
 His Gln His Ile Phe Arg Ala Met Met His Leu Asn Glu Asp Asn Lys
 85 90 95
 Glu Ile Asp Val Val Thr Leu Met Asp Gln Leu Ser Thr Glu Gly Thr
 100 105 110
 Leu Asn Glu Ala Gly Gly Pro Gln Tyr Leu Ala Glu Leu Ser Thr Asn
 115 120 125
 Val Pro Thr Thr Arg Asn Val Gln Tyr Tyr Thr Asp Ile Xaa Ser Xaa
 130 135 140
 Ala Cys Ile Lys Xaa Val Gly Leu Ile Gln Thr Ala Asp Ser Ile Ala
 145 150 155 160

Asn Asp Gly Tyr Xaa Asp Glu Leu Gly Leu Asp Ala Ile Xaa Met Gln
 165 170 175

Asn Val Xaa Phe
 180

<210> 5
 <211> 24
 <212> DNA
 <213> Staphylococcus aureus

<400> 5
 aaattagata aagaagttga aggt 24

<210> 6
 <211> 24
 <212> DNA
 <213> Staphylococcus aureus

<400> 6
 cgagatgatt attgtacgga aaaa 24

<210> 7
 <211> 32
 <212> DNA
 <213> Staphylococcus aureus

<400> 7
 agcatcatcc catggataga atgtatgagc aa 32

<210> 8
 <211> 34
 <212> DNA
 <213> Staphylococcus aureus

<400> 8
 atctattcct cgagcatcat atcagcatgt gcat 34

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/05286**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(G) :C07H 21/00; C12N 1/21, 15/31, 15/63

US CL :435/69.1, 320.1, 252.3; 536/23.7

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 320.1, 252.3; 536/23.7

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

GenBank, EMBL, Swiss-Prot, Pir, Gensearch
search terms: SEQ ID NOS: 1-4**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	NGO et al. Computational complexity, protein structure prediction, and the Levinthal paradox. In The Protein Folding Problem and Tertiary Structure Prediction. Edited by Merz et al. Boston: Birkhauser Boston, 1994, pp. 433, 492-495.	11, 12
A	RUDINGER, J. Characteristics of the amino acid as components of a peptide hormone sequence. In Peptide Hormones. Edited by J.A. Parsons. Baltimore, MD: University Park Press. pp. 1-7.	11, 12
X	WO 97/30070 A1 (SMITHKLINE BEECHAM CORPORATION) 21 August 1997, pages 9, 50, 51, 280, 660, 661.	26, 27

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 MAY 1999

Date of mailing of the international search report

14 JUN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SCOTT D. PRIEBE

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/05286

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	Database Genbank, US National Library of Medicine, (Bethesda, MD, USA), No. D26185, OGASAWARA, N. 'B. subtilis DNA, 180 kilobase region of replication origin'. partial record, 06 February 1999.	1-12, 26, 27

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/05286

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-12, 26, 27

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/05286

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-12, 26, 27, drawn to polynucleotides and first method of using same, making a polypeptide.

Group II, claim(s) 13, 14, 17, drawn to polypeptides and first method of using same, therapy.

Group III, claim(s) 15, 19(b), drawn to antibodies to the polypeptide, and a method of using same.

Group IV, claim(s) 16, 18, drawn to antagonist of polypeptide and method of using same.

Group V, claim(s) 19(a), drawn to second method of using the polynucleotide, sequencing in diagnostic method.

Group VI, claim(s) 20, drawn to second method of using polypeptide, ligand binding assay.

Group VII, claim(s) 21, drawn to third method of using polypeptide, peptide vaccine.

Group VIII, claim(s) 22, drawn to third method of using polynucleotide, DNA vaccine.

Group IX, claim(s) 23-25, drawn to computer readable medium with polynucleotide and amino acid sequences recorded and method for using same.

The inventions listed as Groups I-IX do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: 37 CFR 1.475 has no provision for unity of invention between different, distinct products or between multiple different, distinct methods of using a product. With respect to the polynucleotides and polypeptides, the scope of the polynucleotides claimed is not commensurate with scope of polypeptides claimed in that the polynucleotides include a large fraction that would not encode a polypeptide as claimed.